

Effects of Type II diabetes on capillary hemodynamics in skeletal muscle

Danielle J. Padilla,¹ Paul McDonough,² Brad J. Behnke,³ Yutaka Kano,⁴
K. Sue Hageman,¹ Timothy I. Musch,¹ and David C. Poole¹

¹Departments of Anatomy and Physiology, and Kinesiology, Kansas State University, Manhattan, Kansas;

²Department of Internal Medicine, University of Texas Southwestern Medical Center, Dallas, Texas;

³Division of Exercise Physiology, West Virginia University School of Medicine, Morgantown, West Virginia;

and ⁴Department of Applied Physics and Chemistry, University of Electro-Communications, Chofu, Tokyo, Japan

Submitted 20 March 2006; accepted in final form 5 July 2006

Padilla, Danielle J., Paul McDonough, Brad J. Behnke, Yutaka Kano, K. Sue Hageman, Timothy I. Musch, and David C. Poole.

Effects of Type II diabetes on capillary hemodynamics in skeletal muscle. *Am J Physiol Heart Circ Physiol* 291: H2439–H2444, 2006. First published July 14, 2006; doi:10.1152/ajpheart.00290.2006.—Microcirculatory red blood cell (RBC) hemodynamics are impaired within skeletal muscle of Type I diabetic rats (Kindig CA, Sexton WL, Fedde MR, and Poole DC. *Respir Physiol* 111: 163–175, 1998). Whether muscle microcirculatory dysfunction occurs in Type II diabetes, the more prevalent form of the disease, is unknown. We hypothesized that Type II diabetes would reduce the proportion of capillaries supporting continuous RBC flow and RBC hemodynamics within the spinotrapezius muscle of the Goto-Kakizaki Type II diabetic rat (GK). With the use of intravital microscopy, muscle capillary diameter (d_c), capillary lineal density, capillary tube hematocrit (Hct_{cap}), RBC flux (F_{RBC}), and velocity (V_{RBC}) were measured in healthy male Wistar (control: $n = 5$, blood glucose, 105 ± 5 mg/dl) and male GK ($n = 7$, blood glucose, 263 ± 34 mg/dl) rats under resting conditions. Mean arterial pressure did not differ between groups ($P > 0.05$). Sarcomere length was set to a physiological length ($\sim 2.7 \mu\text{m}$) to ensure that muscle stretching did not alter capillary hemodynamics; d_c was not different between control and GK rats ($P > 0.05$), but the percentage of RBC-perfused capillaries (control: 93 ± 3 ; GK: 66 ± 5 %), Hct_{cap} , V_{RBC} , F_{RBC} , and O_2 delivery per unit of muscle were all decreased in GK rats ($P < 0.05$). This study indicates that Type II diabetes reduces both convective O_2 delivery and diffusive O_2 transport properties within muscle microcirculation. If these microcirculatory deficits are present during exercise, it may provide a basis for the reduced O_2 exchange characteristic of Type II diabetic patients.

intravital microscopy; capillary network; Goto-Kakizaki rat; spinotrapezius; red blood cell flux

TYPE II DIABETES has been associated with decreased exercise tolerance, as demonstrated by slowed oxygen uptake ($\dot{V}O_2$) kinetics at exercise onset (41), lowered maximal $\dot{V}O_2$ ($\dot{V}O_{2\max}$) (4, 43), and a reduction in fractional O_2 extraction (4). Diabetic patients also exhibit compromised muscle blood flow at rest (42) and during exercise (24). The attenuated blood flow may be the result of a blunted endothelium-dependent vasodilation (24, 34, 54) and increased plasma concentrations of the vasoconstrictor endothelin-1 (48). Other changes that accompany this disease include a reduction in capillary density (28, 30) and decreased mitochondrial volume (45) and function (20). This group of patients has also been shown to demonstrate

higher amounts of Type IIb (highly glycolytic) muscle fibers by some (28) but not all (2) investigations.

Recently, preliminary work in our laboratory (37) has shown that muscle microvascular O_2 partial pressures ($PO_{2,mv}$) are reduced by ~ 11 Torr at rest and during contractions in the mixed-fiber type spinotrapezius muscle of the Type II diabetic Goto-Kakizaki (GK) rats (i.e., $PO_{2,mv}$ in GK: ~ 18 Torr vs. control: ~ 29 Torr). This GK model has been considered to be highly representative of the Type II diabetic state in humans (16). The decreased $PO_{2,mv}$ found in the spinotrapezius muscle may be indicative of an impaired microvascular O_2 delivery ($\dot{Q}O_2$) relative to O_2 utilization ($\dot{V}O_2$) (Refs. 6, 32). We reasoned that direct observation of the microcirculation may provide a mechanistic basis for this decreased $PO_{2,mv}$ found in GK rat muscle and offer putative insights into the exercise intolerance typical of diabetic humans (41).

Within skeletal muscle, a functional microvascular bed is necessary for the provision of an adequate supply of O_2 and other nutrients, as well as for removal of metabolic waste products. The modeling studies of Federspiel and Popel (14) suggest that the number of red blood cells (RBCs) adjacent to a muscle fiber (i.e., capillary tube hematocrit multiplied by the length of capillaries) is critical for O_2 exchange, and as the number of RBCs in the capillaries apposed to the myocyte increases, muscle O_2 diffusing capacity is further augmented. Therefore, any structural or functional impairment within the capillary network of skeletal muscle caused by disease that reduces RBC number within the flowing capillaries will negatively impact blood-myocyte O_2 exchange and consequently may play a role in fatigue during repetitive activities. Thus, using intravital microscopy to examine the microcirculation of the spinotrapezius muscle of the GK rat, we hypothesized that the percentage of flowing capillaries would be decreased and that capillary RBC hemodynamics would be impaired in the GK rat. If present, these microcirculatory changes would be expected to attenuate O_2 diffusing capacity, and thus O_2 flux into the myocyte, which would provide one potential mechanism for poor skeletal muscle performance in Type II diabetic patients.

METHODS

Experimental animals. Healthy male Wistar rats (control; $n = 5$; body wt = 557 ± 19 g, 6–8 mo old) and age-matched male GK (Taconic Farm, Germantown, NY) spontaneously diabetic rats ($n = 7$; body wt = 426 ± 15 g) were used in this investigation. The GK rat

Address for reprint requests and other correspondence: D. C. Poole, Dept. of Anatomy/Physiology, College of Veterinary Medicine, 228 Coles Hall, 1600 Denison Ave., Manhattan, KS 66506-5802 (e-mail: poole@vet.ksu.edu).

The costs of publication of this article were defrayed in part by the payment of page charges. The article must therefore be hereby marked “advertisement” in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

is a nonobese, hyperglycemic, insulin-resistant rat strain that was developed by selectively breeding glucose-intolerant Wistar rats (i.e., ~5 generations; Ref. 16). Furthermore, these rats have been reported to have similar or higher non-fasting insulin concentrations compared with age-matched control Wistar rats (36). Goto et al. (17) reported that GK rats require no specially formulated diet and healthy Wistar rats may serve as appropriate controls because they are of the same original strain as the GK rat.

All rats were kept in a controlled environment with a fixed 12:12-h light-dark cycle and with a room temperature maintained at ~22°C. Both control and GK rats were provided conventional rodent chow and water ad libitum. All experimental conditions and surgical procedures were approved by the Kansas State University Institutional Animal Care and Use Committee.

Surgical preparation. Animals were anesthetized with pentobarbital sodium (50 mg/kg ip to effect and supplemented as necessary). The rat was then placed on a heating pad (38°C) to maintain body temperature throughout the experimental protocol. To monitor arterial blood pressure and heart rate (model 200, Digimed BPA, Louisville, KY), the left carotid artery was cannulated (polyethylene-50, Intra-Medic polyethylene tubing, Clay Adams Brands; Sparks, MD). An arterial blood sample was taken from the same catheter during fasting conditions (~6 h fasting) for analysis of blood glucose concentration (Accu-Check Advantage, Roche Diagnostics, Indianapolis, IN) and for measurement of systemic hematocrit (Nova Stat Profile, Waltham, MA).

The spinotrapezius muscle, which lies in the middorsal region of the rat, originates from the lower thoracic and upper lumbar region and inserts onto the spine of the scapula. It is an excellent muscle for intravital studies of the microcirculation because 1) it can be exteriorized without neural or substantial vascular disruption (3, 40); 2) exteriorization permits transmission light microscopy for clear visualization of capillary structures and hemodynamics; 3) a physiological sarcomere length can be set, preventing overstretching of the muscle and associated capillaries, thus avoiding adverse affects to the microcirculation (40); and 4) it is a postural muscle comprising a mixture of fiber types (41% type I, 7% IIa, 17% II/d/x, 35% IIb; Ref. 11) and oxidative capacity similar to human muscle (Ref. 25).

For the experimental preparation, the left spinotrapezius muscle was exteriorized and prepared in situ to examine the microcirculation, as described previously in our laboratory (21, 23, 40, 44, 47). Fascial removal and disturbance was minimized to avoid any associated muscle damage (31). All exposed surrounding tissue, as well as the dorsal surface of the spinotrapezius, was superfused continuously with a Krebs-Henseleit bicarbonate-buffered solution (equilibrated with 95% N₂-5% CO₂, pH 7.4, 38°C). The muscle was sutured (6-0 silk, Ethicon, Somerville, NJ) at five equidistant points around the perimeter to a thin wire horseshoe-shaped manifold, and the exposed dorsal surface of the muscle was protected with Saran Wrap (Dow Brands, Indianapolis, IN) to prevent dehydration until analysis.

The rat was placed on a water circulation-heated Lucite platform, and the spinotrapezius was positioned such that a microvascular field, midway between arteriolar and venular ends within the midcaudal dorsal surface, could be observed using an intravital microscope (Nikon, Eclipse E600-FN; ×40 objective; 0.8 numerical aperture) equipped with a noncontact, illuminated lens, and a high-resolution color monitor (total viewing area = 270 × 210 μm; Sony Trinitron PVM-1954Q, Ichinoya, Japan). The muscle was transilluminated in a fashion that ensured clear resolution of the A-bands of the sarcomeres within one-third to two-thirds of the muscle fibers. The final magnification (×1,184) was confirmed by initial calibration of the system with a stage micrometer (MA285, Meiji Techno). This magnification is adequate for measuring all essential structural and hemodynamic variables (40). The spinotrapezius was maintained at physiological sarcomere length (~2.7 μm) throughout the subsequent observation period, and any exposed tissue was continuously superfused with the Krebs-Henseleit solution.

Experimental design. Once the spinotrapezius muscle was positioned on the platform, 8–10 microvascular viewing fields were each recorded for ~1–1.5 min for every animal. Images were time referenced by frame and fields and stored on Super-VHS high-resolution videocassettes (JVC S-Master XG) by using a videocassette recorder (JVC BR-S822U, Elmwood Park, NJ) for subsequent offline analysis. Mean arterial pressure (MAP) was monitored continuously throughout the data-acquisition period, and the experimental protocol was no longer than 1.5–2 h in duration.

Capillary and fiber structural data analysis. Five of the fields were chosen randomly from each rat for analysis on the basis of clear visualization of sarcomeres, fibers, and capillaries. Capillaries supporting RBC flow were assessed in real time, and each capillary was placed into one of two categories: 1) normal flow = 60 s of continuous, or 2) impeded flow or stopped flow for >10 of 60 s. These criteria were further used for determination of percentage of flowing capillaries [i.e., (number of capillaries supporting RBC flow/total number of visible capillaries per area) × 100]. The presence and direction of RBC flow or the presence of stationary RBCs was also used to determine capillary lineal density (i.e., the number of capillaries per unit muscle width) and countercurrent flow. For all capillaries in which hemodynamics were assessed and where the capillary endothelium was clearly visible on both sides of the lumen, capillary luminal diameter (d_c) was measured (2–4 measurements/capillary) with calipers accurate to ±0.25 mm (±0.17 μm at ×1,184 magnification).

Examination of the microvascular fields was conducted in real time and by frame-by-frame analysis techniques (30 frames/s). Sarcomere length was determined from sets of 10 consecutive in-register sarcomeres (i.e., distance between 11 consecutive A bands) measured parallel to the muscle fiber longitudinal axis. This measurement was repeated 3–4 times where sarcomeres were visible to obtain a mean sarcomere length for each viewing field. For each muscle fiber in which both sarcolemmal boundaries were visible on the screen, the apparent fiber width perpendicular to the longitudinal muscle fiber axis was measured at three locations, and a mean fiber width was determined for each fiber. RBC velocity (V_{RBC}) was determined in all capillaries that were continuously RBC perfused by following the RBC path length over several frames. RBC flux (F_{RBC}) was measured by counting the number of RBCs in a capillary passing an arbitrary point. For each capillary in which hemodynamic data were measured, capillary tube hematocrit (Hct_{cap}) was calculated by the following equation:

$$Hct_{cap} = (\text{volume}_{RBC} \times F_{RBC}) / [\pi \times (d_c/2)^2 \times V_{RBC}]$$

where volume_{RBC} is RBC volume, which was taken to be 61 μm³ (Ref. 1), and capillaries were approximated as circular in cross section (from Ref. 13, as modified by Ref. 23).

Statistical analysis. All data are presented as means ± SE where the group mean is that of the individual muscles rather than individual capillary measurements across muscles. Differences between control and GK groups were tested with a Student's *t*-test. Where there was clear precedence for an a priori directional hypothesis (i.e., decreased F_{RBC} and V_{RBC}), a one-tailed test was used. Statistical significance was accepted at the $P < 0.05$ level.

RESULTS

GK rats exhibited significantly higher fasting blood glucose levels compared with the healthy control rats (control: 105 ± 5 mg/dl; GK: 263 ± 34 mg/dl; $P < 0.05$). The GK rat is considered a nonobese model of Type II diabetes, and this was reflected in the average body weights (control: 557 ± 19 g; GK: 426 ± 15 g; $P < 0.05$). Spinotrapezius weights were also lower in the GK rat (control: 546 ± 25 mg; GK: 421 ± 14 mg; $P < 0.05$); however, when expressed as a ratio to body weight,

Table 1. Cardiovascular variables

Variable	Control (n = 5)	Diabetic (n = 7)
Heart rate, beats/min	371 ± 13	437 ± 14*
Mean arterial pressure, Torr	117 ± 6	110 ± 10
Hematocrit, %	43 ± 1	42 ± 1

Values are means ± SE. * $P < 0.05$

the GK and control groups did not differ ($P > 0.05$). Although sustained within a normal physiological range, heart rate was higher ($P < 0.05$; Table 1) in the GK rats, but MAP did not differ between the groups ($P > 0.05$; Table 1). Systemic (arterial blood) hematocrit, measured at the end of the experiment, did not differ between groups (control: $43 \pm 1\%$; GK: $42 \pm 1\%$; $P > 0.05$; Table 1).

Muscle structural data. Neither sarcomere length (control: $2.6 \pm 0.1 \mu\text{m}$; GK: $2.7 \pm 0.1 \mu\text{m}$) nor capillary diameter (control: $4.7 \pm 0.1 \mu\text{m}$; GK: $5.1 \pm 0.2 \mu\text{m}$) differed between groups (both $P > 0.05$). Although muscle fiber width was lower in the GK rats (control: $65.0 \pm 5.2 \mu\text{m}$; GK: $51.5 \pm 3.9 \mu\text{m}$; $P < 0.05$), total capillary lineal density (i.e., the total number of both perfused and non-RBC-perfused capillaries per unit fiber width) did not differ ($P > 0.05$) between the GK and control rats, but the lineal density of RBC flowing capillaries was attenuated in the GK rats ($P < 0.05$; Fig. 1).

Hemodynamic comparisons. There was a significant decrease in the percentage of RBC flowing capillaries in the GK rats compared with the control rats (control: $93 \pm 3\%$; GK: $66 \pm 5\%$) that occurred in the absence of any alterations in the incidence of countercurrent flowing capillaries (control: $24 \pm 5\%$; GK: $19 \pm 3\%$; $P > 0.05$). In the RBC-perfused capillaries of the individual muscles of GK rats, dramatic decreases were also found in Hct_{cap} (control: $33 \pm 1\%$; GK: $23 \pm 1\%$), V_{RBC} (control: $454 \pm 20 \mu\text{m/s}$; GK: $158 \pm 26 \mu\text{m/s}$), and F_{RBC} (control: $42 \pm 4 \text{RBCs/s}$; GK: $14 \pm 3 \text{RBCs/s}$). Furthermore, V_{RBC} and F_{RBC} were significantly correlated in the individual muscles (r^2 value = 0.928; $P < 0.05$; Fig. 2). Importantly, when the product of flowing lineal density and capillary F_{RBC} was calculated (i.e., an index of \dot{Q}_{O_2}), the blood flow per unit




Fig. 1. Total [i.e., perfused and non-red blood cell (RBC)-perfused capillaries] and flowing (only RBC-perfused capillaries) capillary lineal density in control and Goto-Kakizaki Type II diabetic (GK) rats. Total capillary lineal density did not differ between the 2 groups of animals, but flowing capillary lineal density was significantly lower in the diabetic rats (* $P < 0.05$).




Fig. 2. Correlation between RBC velocity and flux in capillaries supporting RBC flow within spinotrapezius muscles of control and diabetic rats. The average value for all capillaries within a single animal is represented by each data point.

of muscle was markedly lower in the GK rat spinotrapezius compared with control (control: $813 \pm 88 \text{RBCs} \cdot \text{s}^{-1} \cdot \text{mm muscle}^{-1}$; GK: $227 \pm 37 \text{RBCs} \cdot \text{s}^{-1} \cdot \text{mm muscle}^{-1}$; $P < 0.05$; Fig. 3).

DISCUSSION

To the best of our knowledge, this is the first investigation to examine resting skeletal muscle capillary hemodynamics in Type II diabetes, and the data demonstrate that this disease impairs indexes of both perfusive (i.e., \dot{Q}_{O_2}) and diffusive O_2 conductance in the spinotrapezius muscle. Specifically, significant reductions were found in 1) the proportion of continuously RBC-perfused capillaries ($\downarrow 29\%$), 2) capillary V_{RBC} ($\downarrow 65\%$), 3) capillary F_{RBC} ($\downarrow 66\%$), and 4) Hct_{cap} ($\downarrow 30\%$). From these data, an overall reduction in \dot{Q}_{O_2} close to 70% was calculated. These conclusions are consistent with preliminary findings of a reduced $\text{PO}_{2,\text{mv}}$ in the spinotrapezius muscle of GK rats (37) and indicate the presence of a substantial impairment in muscle microcirculatory function in Type II diabetes.

Comparison with published literature. Spinotrapezius muscle fiber widths range from ~ 30 to $60 \mu\text{m}$, with the smallest fibers noted in Type I diabetes (23) and chronic heart failure (44, 47) and the larger fibers (i.e., $60 \mu\text{m}$) found within healthy muscles (23, 44, 47). Within the current study, muscle fiber widths and spinotrapezius weights were significantly smaller in




Fig. 3. The estimated O_2 delivery (\dot{Q}_{O_2} , as calculated by the product of flowing lineal density and RBC flux) was significantly lower in diabetic animals compared with control animals (* $P < 0.05$).

the GK rat versus the control rat, which may have been a potential consequence of the retarded growth reported within the GK rat strain (16). However, when expressed as a ratio of spinotrapezius weight to body weight, the groups did not differ.

In general, the hemodynamic measurements for the control rats in the present investigation are similar to those presented previously for the young healthy rat spinotrapezius (19, 21, 22, 23, 44, 47). This was true despite the body mass being greater in the present investigation (i.e., present, ~560 g; previous, 200–300 g). For example, the percentage of RBC-perfused capillaries (93%) found herein is encompassed within the 80–96% range established for this muscle. In contrast to the above, V_{RBC} and F_{RBC} were both somewhat higher than reported previously, and this may be attributed to some combination of increased age, body mass, and different breed. It is pertinent that Russell and colleagues (47) found an increased V_{RBC} and F_{RBC} in the spinotrapezius of aged (26–28 mo) Fischer \times Brown Norway hybrid rats versus their younger counterparts. In many respects, the capillary hemodynamics of the GK rats (\downarrow %RBC-flowing capillaries, $\downarrow V_{RBC}$, and $\downarrow F_{RBC}$) resemble those found in rats suffering from Type I diabetes (23) or chronic heart failure (21, 44) with the magnitude of these perturbations large enough to discriminate the Type II diabetic rat from any of the healthy populations examined in the studies cited above. One exception to this statement was the Hct_{cap} in the GK animals ($23 \pm 1\%$), which fell within the range established for healthy rats (e.g., Refs. 19, 47) but which was significantly lower than that found in the healthy control rats in the present investigation ($33 \pm 1\%$).

Theoretical basis for hemodynamic alterations. In the present investigation, great care was taken to ensure that the muscles were not stretched to sarcomere lengths that cause a stretch-induced reduction of the capillary luminal diameter and active arteriolar vasoconstriction, both of which would be expected to decrease V_{RBC} and F_{RBC} (22, 39, 40, 52). In contrast, the following mechanisms might be implicated in the hemodynamic impairments found in the GK rats. 1) Increased glycosylation of the RBC membrane protein associated with Type II diabetes would increase RBC rigidity and blood viscosity (9, 27, 33), thereby limiting the ability of RBCs to travel freely through the capillary bed. 2) Acute hyperglycemia (but not chronic) has been reported to adversely affect the microvasculature by altering endothelial glycocalyx function and decreasing functional capillary density (55). This suggests that, preceding the diabetic condition, hyperglycemia compromises endothelial and smooth muscle function and may reduce RBC flux and velocity at the capillary level. 3) Because blood flow to the capillary is primarily controlled at the arteriolar level (Refs. 12, 38), it is quite plausible that the reduced number of RBC-flowing capillaries and slowed capillary hemodynamics may be the result of impaired vasomotor control. The GK rat model does demonstrate a high degree of arteriolar tone consequent to impaired endothelium-dependent vasodilation (7). This phenomenon also occurs in human diabetic patients (15, 33, 49, 51, 54). 4) Similar to that observed in humans (29, 35), GK rats demonstrate increased plasma concentrations of the potent vasoconstrictor endothelin-1 (5). In human diabetic patients, decreased vascular endothelial function has been implicated in the reduced basal forearm blood flow (42, 53) and leg blood flow (24) at rest and during muscle contractions. Thus, from this evidence, there is support for the

notion that vascular dysfunction (endothelial and smooth muscle) in the GK rat may be responsible for the impaired capillary hemodynamics reported in the present investigation. There was certainly no evidence of a reduced capillary luminal diameter (as found in Type I diabetes; Ref. 23) or any structural impediments within the non-RBC-flowing capillaries.

Implications of impaired hemodynamics. According to the elegant modeling studies of Federspiel and Popel (14) and Groebe and Thews (18), the effective O_2 diffusing capacity of muscle ($DO_{2,m}$) is dependent on capillary tube hematocrit (i.e., the number of RBCs contained per unit length of capillary) and the length of RBC-perfused capillaries adjacent to the muscle fibers, as these indexes determine the number of RBCs available for blood-myocyte O_2 transfer at any given instant. Fractional O_2 extraction and thus $PO_{2,mv}$, which constitutes the driving pressure facilitating blood-myocyte O_2 movement, are determined by the relationship between $DO_{2,m}$ and blood flow (\dot{Q}) such that $\dot{V}O_2 = \dot{Q}O_2(1 - e^{-DO_2/\beta\dot{Q}})$ and therefore $\dot{V}O_2/\dot{Q}O_2 = O_2 \text{ extraction} = 1 - e^{-DO_2/\beta\dot{Q}}$, where β is the slope of the O_2 dissociation curve in the physiologically relevant range (46, 50). In the present investigation, we found that Hct_{cap} was significantly decreased in diabetic muscle along with a modest reduction in the lineal density of RBC-perfused capillaries, indicating that $DO_{2,m}$ is expected to be ~40% lower in the GK rat spinotrapezius. With respect to $\dot{Q}O_2$, this index of perfusive O_2 delivery fell ~70% such that the critical ratio $DO_2/\beta\dot{Q}$ and thus O_2 extraction will actually be higher in the GK muscle microcirculation. This finding provides a mechanistic basis for the lowered $PO_{2,mv}$ (37) and greater perturbation of the energetic state (i.e., $\Delta[ADP_{free}]$, $\Delta[\text{phosphocreatine}]$; Ref. 8) observed in muscle of individuals suffering from Type II diabetes.

The impaired capillary hemodynamics evidenced in the Type II diabetic GK rat herein help to explain the reduced $PO_{2,mv}$ (i.e., ~11 Torr) observed recently in this preparation (37) and are also consistent with the reduced limb or muscle blood flows reported in diabetic humans (24, 42, 53). However, at first glance, it is difficult to reconcile a lowered intramuscular $PO_{2,mv}$ (i.e., increased fractional O_2 extraction) with the reduced whole body fractional O_2 extraction reported by Baldi and colleagues (4). One putative explanation is that there is a mismatching of $\dot{Q}O_2$ and $\dot{V}O_2$ within and/or among muscles such that there are regions of under- and overperfusion. Akin to ventilation-perfusion mismatching in the lung (but directionally opposite), the slope of the O_2 dissociation curve at the low $PO_{2,s}$ will dictate that such mismatching acts to decrease overall O_2 extraction and increases venous PO_2 . Specifically, in regions of the very flat portion of the O_2 dissociation curve at low $PO_{2,s}$ (low $\dot{Q}O_2/\dot{V}O_2$), relatively little O_2 is offloaded for additional decrements in PO_2 . In contrast, regions of high $\dot{Q}O_2/\dot{V}O_2$ lie on the steep portion of curve, and maintenance of higher $PO_{2,s}$ translate to substantial elevations of O_2 content. Downstream, when blood from both regions is mixed, the resulting O_2 content will be higher (i.e., fractional O_2 extraction reduced) compared with that draining tissue where there is effective, or at least better, $\dot{Q}O_2/\dot{V}O_2$ matching. Given the pernicious effects of Type II diabetes on vascular control listed above (see *Theoretical basis for hemodynamic alterations*), it is not surprising that such $\dot{Q}O_2/\dot{V}O_2$ mismatch might occur in this disease.

The consequences of a reduced $PO_{2,mv}$ in the diabetic resting muscle may assume greater significance during exercise when greater $\dot{V}O_{2S}$ are required to support cellular energetics. To achieve a given $\dot{V}O_2$ while exercising under conditions of reduced or impaired $\dot{Q}O_2$, diabetic muscle would have to increase fractional O_2 extraction further and exacerbate the decrease in intracellular PO_2 to a level below that found in healthy muscle. As stated above, this situation will reduce the energy state of the myocytes, and this will result in enhanced utilization of glucose, glycogen degradation, and exacerbation of intracellular acid-base disturbances. These events are even more unfavorable when pathological components of a disease such as Type II diabetes includes mitochondrial dysfunction (20, 26) along with impaired glucose uptake and regulation.

Methodological considerations. When interpreting the results of intravital microscopy observations, three concerns are paramount. 1) The exteriorization procedure itself must not impact the measurements themselves. We have demonstrated that neither the spinotrapezius blood flow nor microvascular oxygenation is altered significantly by the surgical interventions necessary to view the microcirculation (3). 2) Image clarity may obscure key structures and bias measurements. It is true that small, non-RBC-perfused capillaries may not be readily discernable. However, Damon and Duling (10) reported that only ~2% of capillaries fell into this category, and those that did would be ineffective for delivering O_2 to the tissue. Notwithstanding this latter point, total capillary density was not different between control and GK muscles, and the principal differences between groups were found in the proportion of RBC-perfused capillaries and their dynamics. 3) Only a relatively small area ($270 \times 210 \mu m$) of tissue per screen was available for observation. Accordingly, the analyses were conducted on five different areas per muscle, and these were not significantly different from one another with respect to the measurements of interest. This finding suggests that the analysis did provide an adequate representation of the microcirculatory hemodynamics occurring in the control and GK spinotrapezius muscles.

In conclusion, the present investigation demonstrates that, within the spinotrapezius muscle of Type II diabetic rats, there is a significant attenuation in the percentage of capillaries supporting RBC perfusion. Moreover, within flowing capillaries, RBC hemodynamics are impaired ($\downarrow V_{RBC}$ and $\downarrow F_{RBC}$), and Hct_{cap} is reduced. These changes occur in the absence of marked structural alterations. If these effects are present during muscle contractions, they may contribute to the O_2 exchange impairment and exercise intolerance characteristic of Type II diabetic patients. Irrespective of this, the GK rat model of Type II diabetes appears to offer a unique window through which to investigate the microcirculatory perturbations that accompany this all-too-prevalent disease. Specifically, the GK rat may prove invaluable for developing and determining the efficacy of different therapeutic modalities designed to limit or reverse the microcirculatory consequences of this disease. One key "next" experiment would be to establish whether correction of blood glucose can reverse the microcirculatory dysfunction established herein.

GRANTS

This work was supported in part by National Institutes of Health Grants HL-69739, HL-67619, HL-50306, and AG-19228.

REFERENCES

- Altman PL and Dittmer DS. *Biology Data Book* (2nd ed.). Bethesda, MD: FASEB, 1974, p. 1598–1613.
- Andersen PH, Lund S, Vestergaard H, Junker S, Kahn BB, and Pedersen O. Expression of the major insulin regulatable glucose transporter (GLUT4) in skeletal muscle of non-insulin-dependent diabetic patients and healthy subjects before and after insulin infusion. *J Clin Endocrinol Metab* 77: 27–32, 1993.
- Bailey JK, Kindig CA, Behnke BJ, Musch TI, Schmid-Schoenbein GW, and Poole DC. Spinotrapezius muscle microcirculatory function: effects of surgical exteriorization. *Am J Physiol Heart Circ Physiol* 279: H3131–H3137, 2000.
- Baldi JC, Aoina JL, Oxenham HC, Bagg W, and Doughty RN. Reduced exercise arteriovenous O_2 difference in Type 2 diabetes. *J Appl Physiol* 94: 1033–1038, 2003.
- Balsiger B, Rickenbacher A, Boden PJ, Biecker E, Tsui J, Dashwood M, Reichen J, and Shaw SG. Endothelin A-receptor blockade in experimental diabetes improves glucose balance and gastrointestinal function. *Clin Sci (Lond)* 103, Suppl 48: 430S–433S, 2002.
- Behnke BJ, Barstow TJ, Kindig CA, McDonough P, Musch TI, and Poole DC. Dynamics of oxygen uptake following exercise onset in rat skeletal muscle. *Respir Physiol Neurobiol* 133: 229–239, 2002.
- Bitar MS, Wahid S, Mustafa S, Al-Saleh E, Dhaunsi GS, and Al-Mulla F. Nitric oxide dynamics and endothelial dysfunction in type II model of genetic diabetes. *Eur J Pharmacol* 511: 53–64, 2005.
- Challiss RA, Vranic M, and Radda GK. Bioenergetic changes during contraction and recovery in diabetic rat skeletal muscle. *Am J Physiol Endocrinol Metab* 256: E129–E137, 1989.
- Chung TW, Liu AG, and Yu JH. Increased red cell rigidity might affect retinal capillary blood flow velocity and oxygen transport efficiency in type II diabetes. *Diabetes Res* 23: 75–82, 1993.
- Damon DH and Duling BR. Distribution of capillary blood flow in the microcirculation of the hamster: an in vivo study using epifluorescent microscopy. *Microvasc Res* 27: 81–95, 1984.
- Delp MD and Duan C. Composition and size of Type I, IIA, IID/X, and IIB fibers and citrate synthase activity of rat muscle. *J Appl Physiol* 80: 261–270, 1996.
- Delp MD and Laughlin MH. Regulation of skeletal muscle perfusion during exercise. *Acta Physiol Scand* 162: 411–419, 1998.
- Desjardins C and Duling BR. Heparinase treatment suggests a role for the endothelial cell glycocalyx in regulation of capillary hematocrit. *Am J Physiol Heart Circ Physiol* 258: H647–H654, 1990.
- Federspiel WJ and Popel AS. A theoretical analysis of the particulate nature of blood on oxygen release in capillaries. *Microvasc Res* 32: 164–189, 1986.
- Goodfellow J, Ramsey MW, Luddington LA, Jones CJ, Coates PA, Dunstan F, Lewis MJ, Owens DR, and Henderson AH. Endothelium and inelastic arteries: an early marker of vascular dysfunction in non-insulin dependent diabetes. *Br Med J* 312: 744–745, 1996.
- Goto Y and Kakizaki M. The spontaneous-diabetes rat: a model on non-insulin dependent diabetes mellitus. *Proc Jpn Acad* 57: 381–384, 1981.
- Goto Y, Kakizaki M, and Masaki N. Spontaneous diabetes produced by selective breeding of normal Wistar rats. *Proc Jpn Acad* 51: 80–85, 1975.
- Groebe K and Thews G. Calculated intra- and extracellular PO_2 gradients in heavily working red muscle. *Am J Physiol Heart Circ Physiol* 259: H84–H92, 1990.
- Kano Y, Padilla DJ, Behnke BJ, Hageman KS, Musch TI, and Poole DC. Effects of eccentric exercise on microcirculation and microvascular oxygen pressures in rat spinotrapezius muscle. *J Appl Physiol* 99: 1516–1522, 2005.
- Kelley DE, He J, Menshikova EV, and Ritov VB. Dysfunction of mitochondria in human skeletal muscle in type 2 diabetes. *Diabetes* 51: 2944–2950, 2002.
- Kindig CA, Musch TI, Basaraba RJ, and Poole DC. Impaired capillary hemodynamics in skeletal muscle of rats in chronic heart failure. *J Appl Physiol* 87: 652–660, 1999.
- Kindig CA and Poole DC. Sarcomere length-induced alterations of capillary hemodynamics in rat spinotrapezius muscle: vasoactive vs. passive control. *Microvasc Res* 61: 64–74, 2001.
- Kindig CA, Sexton WL, Fedde MR, and Poole DC. Skeletal muscle microcirculatory structure and hemodynamics in diabetes. *Respir Physiol* 111: 163–175, 1998.

24. Kingwell BA, Formosa M, Muhlmann M, Bradley SJ, and McConell GK. Type 2 diabetic individuals have impaired leg blood flow responses to exercise: role of endothelium-dependent vasodilation. *Diabetes Care* 26: 899–904, 2003.
25. Leek BT, Mudaliar SR, Henry R, Mathieu-Costello O, and Richardson RS. Effect of acute exercise on citrate synthase activity in untrained and trained human skeletal muscle. *Am J Physiol Regul Integr Comp Physiol* 280: R441–R447, 2001.
26. Lowell BB and Shulman GI. Mitochondrial dysfunction and type 2 diabetes. *Science* 307: 384–387, 2005.
27. MacRury SM, Small M, MacCuish AC, and Lowe GD. Association of hypertension with blood viscosity in diabetes. *Diabet Med* 5: 830–834, 1988.
28. Mårin P, Andersson B, Krotkiewski M, and Bjorntorp P. Muscle fiber composition and capillary density in women and men with NIDDM. *Diabetes Care* 17: 382–386, 1994.
29. Mather KJ, Mirzamohammadi B, Lteif A, Steinberg HO, and Baron AD. Endothelin contributes to basal vascular tone and endothelial dysfunction in human obesity and type 2 diabetes. *Diabetes* 51: 3517–3523, 2002.
30. Mathieu-Costello O, Kong A, Ciaraldi TP, Cui L, Ju Y, Chu N, Kim D, Mudaliar S, and Henry RR. Regulation of skeletal muscle morphology in type 2 diabetic subjects by troglitazone and metformin: relationship to glucose disposal. *Metabolism* 52: 540–546, 2003.
31. Mazzoni MC, Skalak TC, and Schmid-Schonbein GW. Effects of skeletal muscle fiber deformation on lymphatic volumes. *Am J Physiol Heart Circ Physiol* 259: H1860–H1868, 1990.
32. McDonough P, Behnke BJ, Kindig CA, and Poole DC. Rat muscle microvascular PO_2 kinetics during the exercise off-transient. *J Exp Biol* 86: 349–356, 2001.
33. McMillan DE. Exercise and diabetic microangiopathy. *Diabetes* 28: S103–S106, 1979.
34. McVeigh GE, Brennan GM, Johnston GD, McDermott BJ, McGrath LT, Henry WR, Andrews JW, and Hayes JR. Impaired endothelium-dependent and independent vasodilation in patients with type 2 (noninsulin-dependent) diabetes mellitus. *Diabetologia* 35: 771–776, 1992.
35. Nugent AG, McGurk C, Hayes JR, and Johnston GD. Impaired vasoconstriction to endothelin 1 in patients with NIDDM. *Diabetes* 45: 105–107, 1996.
36. Ostenson CG. The Goto-Kakizaki rat. In: *Animal Models of Diabetes: A Primer*, edited by Sima AAF and Shafir E. Amsterdam, The Netherlands: Harwood Academic, 2001, p. 197–211.
37. Padilla DJ, McDonough P, Behnke BJ, Kano Y, Hageman KS, Musch TI, and Poole DC. Effects of Type II diabetes on muscle microvascular oxygen pressures (Abstract). *The Physiologist* 47: 351(A35.3), 2004.
38. Pohl U, De Wit C, and Gloe T. Large arterioles in the control of blood flow: role of endothelium-dependent dilation. *Acta Physiol Scand* 168: 505–510, 2000.
39. Poole DC and Mathieu-Costello O. Capillary and fiber geometry in rat diaphragm perfusion fixed in situ at different sarcomere lengths. *J Appl Physiol* 73:151–159, 1997.
40. Poole DC, Musch TI, and Kindig CA. In vivo microvascular structural and functional consequences of muscle length changes. *Am J Physiol Heart Circ Physiol* 272: H2107–H2114, 1997.
41. Regensteiner JG, Bauer TA, Reusch JE, Brandenburg SL, Sippel JM, Vogelsson AM, Smith S, Wolfel EE, Eckel RH, and Hiatt WR. Abnormal oxygen uptake kinetic responses in women with type II diabetes mellitus. *J Appl Physiol* 85: 310–317, 1998.
42. Regensteiner JG, Popylisen S, Bauer TA, Lindenfeld J, Gill E, Smith S, Oliver-Pickett CK, Reusch JEB, and Weil JV. Oral L-arginine and vitamins E and C improve endothelial function in women with type 2 diabetes. *Vasc Med* 8: 169–175, 2003.
43. Regensteiner JG, Sippel J, McFarling ET, Wolfel EE, and Hiatt WR. Effects of non-insulin dependent diabetes on oxygen consumption during treadmill exercise. *Med Sci Sports Exerc* 27: 875–881, 1995.
44. Richardson TE, Kindig CA, Musch TI, and Poole DC. Effects of chronic heart failure on skeletal muscle capillary hemodynamics at rest and during contractions. *J Appl Physiol* 95: 1055–1062, 2003.
45. Ritov VB, Menshikova EV, He J, Ferrell RE, Goodpaster BH, and Kelley DE. Deficiency of subsarcolemmal mitochondria in obesity and type 2 diabetes. *Diabetes* 54:8–14, 2005.
46. Roca J, Agusti AG, Alonso A, Poole DC, Viegas C, Barbera JA, Rodriguez-Roisin R, Ferrer A, and Wagner PD. Effects of training on muscle O_2 transport at $\dot{V}O_{2\max}$. *J Appl Physiol* 73: 1067–1076, 1992.
47. Russell JA, Kindig CA, Behnke BJ, Poole DC, and Musch TI. Effects of aging on capillary geometry and hemodynamics in rat spinotrapezius muscle. *Am J Physiol Heart Circ Physiol* 285: H251–H258, 2003.
48. Schneider JG, Tilly N, Hierl T, Sommer U, Hamann A, Dugi K, Leidig-Bruckner G, and Kasperk C. Elevated plasma endothelin-1 levels in diabetes mellitus. *Am J Hypertens* 15: 967–972, 2002.
49. Steinberg HO, Chaker H, Leaming R, Johnson A, Brechtel G, and Baron AD. Obesity/insulin resistance is associated with endothelial dysfunction. Implications for the syndrome of insulin resistance. *J Clin Invest* 97: 2601–2610, 1996.
50. Wagner PD, Hoppeler H, and Saltin B. Determinants of maximal O_2 uptake. In: *The Lung: Scientific Foundations*, edited by Crystal RG, West JB, Weibel ER, and Barnes PJ. New York: Raven, 1997, p. 2033–2041.
51. Watts GF, O'Brien SF, Silvester W, and Millar JA. Impaired endothelium-dependent and independent dilatation of forearm resistance arteries in men with diet-treated non-insulin-dependent diabetes: role of dyslipidaemia. *Clin Sci (Lond)* 91: 567–573, 1996.
52. Welsh DG and Segal SS. Muscle length directs sympathetic nerve activity and vasomotor tone in resistance vessels of hamster retractor. *Circ Res* 79: 551–559, 1996.
53. Yu HI, Sheu WH, Lai CJ, Lee WJ, and Chen YT. Endothelial dysfunction in type 2 diabetes mellitus subjects with peripheral artery disease. *Int J Cardiol* 78: 19–25, 2001.
54. Williams SB, Cosco JA, Roddy MA, Johnstone MT, and Creager MA. Impaired nitric oxide-mediated vasodilation in patients with non-insulin-dependent diabetes mellitus. *J Am Coll Cardiol* 27: 567–574, 1996.
55. Zuurbier CJ, Demirci C, Koeman A, Vink H, and Ince C. Short-term hyperglycemia increases endothelial glycocalyx permeability and acutely decreases lineal density of capillaries with flowing red blood cells. *J Appl Physiol* 99: 1471–1476, 2005.